INTERNATIONAL AYURVEDIC MEDICAL JOURNAL

IAMJ

Research Article

ISSN: 2320 5091

Impact Factor: 4.018

A COMPARATIVE STUDY OF MAHISHA KSHEERA AND SHUDDHA BALA TAILA DHARA IN NIDRANASHA W.S.R. INSOMNIA

Karthikeya Prasad¹, Gopidev T. G²

¹ Guide; ²PG Scholar;

Department of PG Studies in Panchakarma, Karnataka Ayurveda Medical College and Hospital, Ashoknagara, Mangalore, Karnataka, India

Email: drkarthikshet@gmail.com

ABSTRACT

Sleep or *nidra*, is an indicator of good health, is one of the natural instincts of man. *Nidranasha* is the term commonly used to denote the state of reduced or disturbed sleep. Insomnia can be correlated with *nidranasha*. In *ayurveda doshas* involved in *nidranasha* are *vata* and *pitta*. The management includes *doshasamana* as well as subjective *swasthya*. The objectives of the present study are to evaluate efficacy of *shirodhara* with *mahishaksheera* and *suddhabalataila* in *nidranasha* before and after treatment in the group and between the groups. **Method:** The outcome of the study is evaluated clinically on 40 patients divided into 2 groups A and B each 20 patients. For Group A *shirodhara* with *mahishaksheera* and for Group B *shirodhara* with *suddhabalataila* was done for 45min, for 7 consecutive days. Observations were made on 1st, 7th and 21st day. In both groups objective and subjective parameters were assessed and the results were statistically analysed.

Keywords: nidranasha, insomnia, shirodhara, suddhabalataila, mahishaksheera

INTRODUCTION

Nidra is a state, when manas, jnanendriya, karmendriya get exhausted, distract from the state of objects¹ and thus man goes into a state of sleep. During the description of nidrautpatti, acharyas have mentioned that sharirika dosha kapha manasika dosha tama are responsible for nidra². Ayurveda mentioned nidra as one among the trayopasthambha³ to explain the importance of nidra. Sleep is characterised by a reversible state of relative unresponsiveness and tranquillity. The condition happens almost regularly and repetitively every day. Nearly 1/3rd of one's life span is spent in sleep.

• Nidranasha or loss of sleep (insomnia) is caused by aggravation of vata and pitta, exhaustion of the mind and also due to injury to the body⁴. The inter relationship of vata and manas by saying " chalevatechalam chittam nischale nischalambhavet"⁵ is well said in Hatayoga Pradeepa. The essential feature of insomnia disorder is dissatisfaction with sleep quantitatively or qualitatively and or complaints of difficulty initiating or maintaining sleep. The sleep complaints are accompanied by clinically significant distress or impairment in social, occupational or other important areas of functioning⁶. Statistically, insomnia is global epidemic affecting quality of life of nearly 45% of the world population⁷. Different treatment principles are mentioned in classics like *abhyanga*, *utsadana*, *snana*, *chakshutarpana*, *lepa* on *shira* for inducing sleep⁸. Practically *shirodhara* is considered as the best modality of the available treatment. Insomnia, being an alarming problem, needs safe and effective treatment. Hence the present study is planned.

In Ayurveda, Shirodhara is considered as one of the snehana karma widely practiced for treating vatavyadhis, nidranasha etc⁹. Shirodhara is the process in which medicated potion like oil, milk, or kwatha is poured as a continuous stream on the forehead from a specified distance and then allowed to run over the scalp for a stipulated period of time.

In the present study ShuddaBalaTaila and Mahishaksheera is used. Shuddhabalataila contains Bala, tilataila, ksheera¹⁰ all of which are having vatapittahara properties^{11,12}. Also in the classics shirodhara with ksheera is indicated in cases of nidranasha¹³. Mahishaksheera has madhura rasa, snigdhaguna, sitaveerya and madhura vipaka which subsides vata and pitta doshas and increases the kaphadosha, it also increases tamoguna in the shirahpradesha¹⁴.

With the procedure of Shirodhara, vata and pitta doshas get pacified and there is elevation in kaphadosha. It is also stated that when an individual lies down in a relaxed state for longer time, tamasi-kaguna over-comes the rajasikaguna. Also the procedure of Shirodhara brings the sanjnavahasrotas to a peaceful state of rest which in turn helps in inducing sleep¹⁴.

AIM AND OBJECTIVES:

- To study and evaluate the efficacy of Mahishaksheera Shirodhara in Nidranasha.
- To study and evaluate the efficacy of Shuddha-Bala Taila Shirodhara in Nidranasha.

- To compare and evaluate the efficacy of Mahisha Ksheera Shirodhara and Shuddhabala taila Shirodhara in Nidranasha.

MATERIALS AND METHODS:

SOURCE OF DATA

Patients who were fulfilling the criteria for diagnosis and inclusion were selected from outpatient department of Karnataka Ayurveda medical college,

Mangaluru.

METHODS OF COLLECTION OF DATA

- a) Clinical study was conducted on 40 patients of *nidranasha*.
- b) Diagnostic criteria:

Diagnosis will be made on the basis of the following lakshanas of nidranasha and DSM V criteria for insomnia:

- 1. lakshana of nidranasha
- moha
- angamarda
- akshigaurava
- shirashoola
- alasya
- jrumbha
- tandra
- diagnostic and statistical manual of mental disorders, fifth edition text revision (DSM V criteria for insomnia)
- a predominant complaint of dissatisfaction with sleepy quantity or quality, associated with one (or more) of the following symptoms:
- difficulty initiating sleep
- difficulty maintaining sleep, characterized by frequent awakenings or problems returning to sleep after awakenings
- early morning awakening with inability to return to sleep
- ➤ the sleep difficulty is present for at least 3nights per week
- > the sleep difficulty is present for at least 3 months

Inclusion criteria: c)

- Patients of both sexes, between the age group of
- patients fulfilling the diagnostic criteria of nidranasha

d) Exclusion criteria:

- nidranasha due to other conditions like madatyaya, abhighata
- patients with psychiatric diagnosis are excluded from this study were receiving any medication or drugs that could cause sedation
- patients with other medical diseases like orthostatic hypotension, liver disease, thyroid disease, heart disease, history of seizure or cognitive impairment
- pregnancy
- patient unfit for shirodhara

INTERVENTION:

TREATMENT GROUP

40 patients with a definite diagnosis of fulfilling the diagnostic, inclusion criteria of nidranasha where selected and grouped into 2 groups of size each 20.

GROUP I Shirodhara with MahishaKsheera

: mahishaksheera drug

: 1500ml fresh mahishaksheera per day. quantity

duration : 7 days.

follow up : 14days after treatment

purvakarma : purvakarma is related with the preparation of the patient.

pradhana karma : shirodehara with mahishaksheera duration : dhara procedure is done for 45mint :mahishaksheera is change every day. period for changing the liquid

: it will be lukewarm temp temperature of liquid : samsarjana karma followed paschatkarma

GROUP II Shirodara with ShudhaBalaTaila

: shudhabalataila drug

: 1500ml fresh shudhabalataila per day. quantity

: 7 days. duration

: 14days after treatment follow up

: purvakarma is related with the preparation of the patient. purvakarma

: shirodhara with shudhabalataila pradhana karma : dhara procedure is done for 45 mint duration : maintain 1500ml of taila daily period for changing the liquid : it will be lukewarm temp temperature of liquid : samsarjana karma followed.

ASSESSMENT CRITERIA

paschatkarma

• the improvement in the following signs and symptoms of the disease was assessed by grading of different parameters

CRITERIA FOR ASSESSMENT OF OVERALL **EFFECT**

Overall effect of the therapy and reduction in symptoms was assessed in terms of grading of subjective and objective criteria. It was observed by adopting the following criteria.

GRADING OF THE SUBJECTIVE AND OBJECTIVE CRITERIA:

Sl No	Chief complints	Sec	Score Range (0-3)					
		Ran						
1	Total sleep duration	0	1	2	3			
2	Sleep induction	0	1	2	3			
3	Awakenings during the night	0	1	2	3			
4	Final awakening earlier than desired	0	1	2	3			
5	Overall quality of sleep	0	1	2	3			
6	Sense of wellbeing during the day	0	1	2	3			
7	Functioning (physical and mental) during the day	0	1	2	3			
8	Sleepiness during the day	0	1	2	3			

Sl No	Associated complaints	Sco	Score Range (0-3)				
		Rang					
1	Moha	0	1	2	3		
2	Angamarda	0	1	2	3		
3	Akshigaurav	0	1	2	3		
4	Shirashoola	0	1	2	3		
5	Alasya	0	1	2	3 .		
6	Jrimbha	0	1	2	3		
7	Tandra	0	1	2	3		

(0- No problem 1-Mild 2-Moderate 3-Severe)

OBSERVATION AND RESULT

Effect of therapy on objective parameters between the groups- After treatment

Table 1: Effect of therapy on objective parameters between the groups- After treatment

parameter	N		Mean rank	ζ	Sum of ran	ks	Test	P value
	Group A	Group B	Group A	Group B	Group A	Group B	value	
Total sleep duration	20	20	21.5	19.5	430	390	180	.298
Sleep induction	20	20	20.95	20.5	419	401	191	.782
Awakenings during the night	20	20	20.98	20.03	419.5	400.5	190.5	.742
Final awakening earlier than desired	20	20	23.10	17.90	462	358	148	.096
Overall quality of sleep	20	20	22	19	440	380	170	.343
Sense of wellbeing during the day	20	20	21.50	19.50	430	390	180	.471
Functioning (physical and mental) during the day	20	20	21.50	19.50	430	390	180	.471
Sleepiness during the day	20	20	21.50	19.50	430	390	180	.496

From the data available the following conclusions are made between the groups for objective parameters. The Mann Whitney U test was done as the data

does not follow normality. To know significance of treatment between the groups mean ranks were compared.

Effect of therapy on subjective parameters between the groups- After treatment

Table 2: Effect of therapy on subjective parameters between the groups- After treatment

parameter	N		Mean rank		Sum of ranks		Test	P value
	Group A	Group B	Group A	Group B	Group A	Group B	value	
Moha	20	20	21	20	420	400	190	.637
Angamarda	20	20	21	20	420	400	190	.727
Akshigaurav	20	20	20.5	20.5	410	410	200	.1000
Shirashoola	20	20	21	20	420	400	190	.755
Alasya	20	20	22	19	440	380	170	.262
Jrimbha	20	20	20.5	20.5	410	410	200	1.000
Tandra	20	20	22.5	18.5	450	370	160	.289

Effect of therapy on objective parameters between the groups- After follow-up

Table 3: Effect of therapy on objective parameters between the groups- After follow

parameter	N		Mean rank		Sum of ranks		Test	P value
	Group A	Group B	Group A	Group B	Group A	Group B	value	
Total sleep duration	20	20	22.03	18.98	440.50	379.50	169.5	.211
Sleep induction	20	20	20.45	20.55	409	411	199	.971
Awakenings during the night	20	20	22.30	18.70	446	374	164	.227
Final awakening earlier than desired	20	20	23.55	17.65	467	353	143	.079
Overall quality of sleep	20	20	23.50	17.50	470	350	140	.060
Sense of wellbeing during the day	20	20	24	17	480	340	130	.014
Functioning (physical and mental) during the day	20	20	26	15	520	300	90	.000
Sleepiness during the day	20	20	20	21	420	400	190	.708

From the data available the following conclusions are made between the groups for objective parameters. The Mann Whitney U test was done as the data

does not follow normality. To know significance of treatment between the groups mean ranks were compared.

Effect of therapy on subjective parameters between the groups- After follow-up

Table 4: Effect of therapy on subjective parameters between the groups- After follow

parameter	ter N		Mean rank			nks	Test	P value
•	Group A	Group B	Group A	Group B	Group A	Group B	value	
Moha	20	20	21.5	19.5	430	390	180	.298
Angamarda	20	20	21.5	19.5	430	390	180	.471
Akshigaurav	20	20	20.5	20.5	410	410	200	1.000
Shirashoola	20	20	22.5	18.5	450	370	160	.212
Alasya	20	20	22.5	18.5	450	370	160	.037
Jrimbha	20	20	21.93	19.08	438.5	381.5	171.5	.179
Tandra	20	20	22.5	18.5	450	370	160	.080

DISCUSSION

Probable mode of action of *shirodhara* and drugs on its properties

Shirodhara is a purifying and rejuvenating therapy designed to eliminate toxins and mental exhaustion as well as relieve stress and any ill effects on the central nervous system.

Procedural effect of the process:

To discuss Procedural effect of *Shirodharakarma*, it dived in to three steps effects.

- 1. Penetration
- 2. Stimulation
- 3. Relaxation

1. Penetration:

In mahishiksheeradhara and sudhabalatailadhara on the forehead of the patients, it stuck little bit on fore head which consist some vital centres such as Agya Chakra, Sthapnimarmas. SukshmaIndriyasresides in Mastishkaand Sneha ingredients enter through the penetration in the skin of forehead. The Penetration power is depended on consistency and density of drugs. In this study drug of sudhabalaTaila has greater penetration power than mahishaksheera, as has lipid constitution which can be penetrated easily through cell walls. So, it can be said that sudhabalatailadhara is more beneficial than mahishaksheeradhara.

2. Stimulation

To reach the central nervous system, a drug must have a high degree of lipid solubility or a specialized transport mechanism. After penetration, it can be said it stimulates CNS. Diminished function of thalamus and forebrain causes serotonin levels to decrease and catecholamine's increase thus manifesting insomnia. Shirodhara normalizes the functions of thalamus and forebrain, which brings the amount of serotonin and catecholamine's to normal stage, which induces sleep. Shirodhara stimulates the pineal gland which produces the hormone melatonin, which induced nidrajanana property. According to modern physiology, drugs may act directly on neurons and modify the neuronal functions. They may act by sending afferent impulses to the central

nervous system via the chemoreceptors, baroreceptors and peripheral nerves and there by eliciting psychic, somatic or visceral responses. They may affect the nutrition and oxygen supply of the CNS by altering its blood supply or affecting its metabolism.

3. Relaxation:

In Shirodhara, patients feel relaxation both physically as well as mentally.

Relaxation of the frontalis muscle tends to normalize the entire body and achieve a decrease activity of SNS with lowering of brain cortisone and adrenaline level; synchronizes the brain wave (alpha waves) strengthens the mind and spirit and this continues even after the relaxation. Shirodhara works on cerebral system helps in relaxing the nervous system and balancing the PranaVayu around the head. It improves the function of five senses, helps in insomnia, stress, anxiety, depression, hair loss, fatigue, imbalance of Vata and makes one calm and fresh accompanied by distress or impairment in day time functioning.

CONCLUSION

The main conclusions derived from the study are

- Demo graphic data showed that people of age group. 41-50, females, working in various field, married, middle class are predominant in the study.
- Shirodhara with mahishaksheera significantly effective in nidranasha.
- Shirodhara with sudhabalataila significantly effective in nidranasha
- The results were assessed based on the relief obtained to the subjective and objective parameters in the group and between the groups. Both groups were found to be effective in *nidranasha*, on after treatment (7th day) results there is no much statistical significant difference between groups. In follow up (21st day) Awakening during night, final awakening earlier than desired, *alasya* are statistically significant in group B than group A. This indicates that *shirodhara* with *suddhaba*-

lataila is more effective than shirodhara with mahishaksheera,

LIMITATIONS

- The limited size of sample was not sufficient to conclude the efficacy of the intervention
- · Limited study period
- Lack of willingness of patients to undergo IP treatment.
- EEG co relation and differential diagnosis was not made.
- Study could have been more significant if it screened persons according to their prakrithi.

RECOMMENDATIONS

- · Larger sample size
- Larger period of study
- Compare the effects of shirodhara with other indicated procedure

REFERENCES

- Sharma RK. Agnivesa's caraka samhitha. Reprint ed. Varanasi: Chowkhamba Sanskrit Series Office;2008.vol I .p-381
- Murthy Srikantha KR. Susruta samhitha. Reprint ed. Varanasi: chaukhambha orientalia;2008.vol I.p.69
- Sharma RK. Agnivesa's caraka samhitha. Reprint ed. Varanasi: Chowkhamba Sanskrit Series Office;2008.vol I.p-219
- Murthy Srikantha KR. Susruta samhitha. Reprint ed. Varanasi: chaukhambha orientalia;2008.vol I.p.65
- Swatmarama yogindra, Hatayoga pradeepika, Brahmaranda, Mumbai, khemraj shrikrishnada; dvitiyopadesha, 2006, p-41
- American psychiatry association. Diagnostic and stasticial manual of mental disorders. Washington. DC,londan, England: American psychiatry publishing;5th ed. 2013 p-361
- Seema chawardol G, sapan jain B, clinical evaluation of role of shirodhara and powder of valerianawallichii in the management of insomnia. World journal of pharmacy and pharmaceutical sciences. [serial online] 2014july12 volume-3, issue 9- p-1578,1579 URL: article wjpps 1410169487

- Sharma RK. Agnivesa's caraka samhitha. Reprint ed. Varanasi: Chowkhamba Sanskrit Series Office;2008.vol I.p-384
- Krishnan KV, gopala pillai S. Sahasrayogam Sujanapriya commentary. 30thed. Mullakkal: vidhyarambam publishers;2011.p.472
- Krishnan KV, gopala pillai S. Sahasrayogam Sujanapriya commentary. 30thed. Mullakkal: vidhyarambam publishers;2011.p.312,313
- 11. Sastry J.L.N. Dravya Guna Vijnana Varanasi: Chaukhambha orientalia.vol IIp-89
- Sreekumar T. astanga hridaya. Mannuty: published by, harisree hospital.P.124
- Sharma RK. Agnivesa's caraka samhitha. Reprint ed. Varanasi: Chowkhamba Sanskrit Series Office;2008.vol.p-553
- Pokharel sanjay, Sharma ajay kumar, evaluation of insomrid tablet and shirodhara in the management of anidra. An international quarterly journal of research inayurveda. [serial online] 2010 janmar; 31(1)4047, doi:10.4103/09748520.68209: URL:www.ncbi.nlm.nih.gov/pmc/articles/pmc32153

Source of Support: Nil Conflict Of Interest: None Declared

How to cite this URL: Karthikeya Prasad & Gopidev T.G: A Comparative Study Of Mahisha Ksheera And Shuddha Bala Taila Dhara In Nidranasha W.S.R. Insomnia. International Ayurvedic Medical Journal (online) 2018 (cited September, 2018) Available from: http://www.iamj.in/posts/images/upload/1949 1955.pdf